# 2 Important Areas for New Zealand Seabirds

# Sites on Land

Rivers, estuaries, coastal lagoons & harbours









### IMPORTANT AREAS FOR NEW ZEALAND SEABIRDS



This document has been prepared for Forest & Bird by Chris Gaskin, IBA Project Coordinator (NZ).

The Royal Forest & Bird Protection Society of New Zealand Level One, 90 Ghuznee Street PO Box 631 Wellington 6140 NEW ZEALAND

This report is available from the Forest & Bird website in pdf form.

© Copyright February 2016, Forest & Bird

#### Contributors

The following individuals have contributed to the profiles in this document in a variety of ways, including supply of data and information about seabirds, and reviewing draft material, site profiles, species lists and site maps.

Nick Allen, Tim Barnard, Tony Beauchamp, Mike Bell, Mark Bellingham, Robin Blyth, Phil Bradfield, John Cheyne, Wynston Cooper, Andrew Crossland, Philippa Crisp, Paul Cuming, John Dowding, Hannah Edmonds, Lloyd Esler, Julian Fitter, Peter Frost, Mel Galbraith, Liz Garson, Peter Gaze, Andrew Grant, Tony Habraken, Kate Hand, Ken Hughey, Elaine Lagnaz, Chris Lalas, Peter Langlands, David Lawrie, Eila Lawson, Nick Ledgard, Nikki McArthur, Rachel McClellan, Craig McKenzie, Bruce McKinlay, Michael McSweeney, David Melville, Gary Melville, Mark O'Brien, Colin O'Donnell, Gwenda Pulham, Aalbert Rebergen, Phil Rhodes, Adrien Riegen, Neil Robertson, Paul Sagar, Frances Schmechel, Rob Schuckard, Ian Southey, Kate Steffens, Graeme Taylor, Gillian Vaughan, Jan Walker, Susan Waugh, David Wilson, Kerry-Jayne Wilson, Steve Wood, Keith Woodley.

Cover design: Danielle McBride, Paradigm Associates, Auckland

Front cover: Rachel McLellan (Black-billed Gulls), Craig McKenzie (Black-fronted Tern)

Back cover: Frederic Pelsy (Ahuriri River)

Recommended citation: Forest & Bird (2016). New Zealand Seabirds: Sites on Land, Rivers, estuaries, coastal lagoons & harbours. The Royal Forest & Bird Protection Society of New Zealand, Wellington, New Zealand.

NZ104 Whitestone River Southland	
----------------------------------	--

Location	New Zealand, Southland	
Central coordinates	45.460913°S, 167.788439°E (SH94 bridge)	
IBA criteria	A1	
Area	1,067 ha	
Altitude	255 - 560m	
Year of Assessment	2013	

## Populations of IBA trigger species:

Species	Season	Period	Population	IBA	IUCN	NZ Threat
			estimate	criteria	cat.	Class.
Black-billed Gull	Breeding	1997-1998	1,000 (nests)	A1, A4i	EN	Nationally Critical
Black-fronted Tern	Breeding		No count	A1	EN	Nationally Vulnerable


Ornithological information: Besides the trigger species the following species are confirmed or likely to be breeding: Black Shag, White-faced Heron, Canada Goose, Paradise Shelduck, Mallard, Grey Duck, Australasian Harrier, NZ Falcon, South Island Pied Oystercatcher, Pied Stilt, Banded Dotterel, Spur-winged Plover, Southern Black-backed Gull, Rock Pigeon, Welcome Swallow, NZ Pipit, South Island Fantail, Silvereye, Skylark, Blackbird, Song Thrush, Yellowhammer, Chaffinch, Greenfinch, Goldfinch, Redpoll, House Sparrow, Starling, Australian Magpie.

Protected area	Designation	Relationship with IBA
Whitestone River	Marginal Strips	

The IBA borders and includes sections of the Snowdon Forest Conservation Area.

IUCN Habitat	Habitat detail	Extent (% of site)
	Permanent Rivers; Extensive River Gravel Beds	Major
Introduced Vegetation	Broom, gorse, pasture grasses & rank weeds, willow margins	Major
Grasslands	Tussock on river flats (upper reaches)	Major
Shrubland	Manuka, matagouri (Discaria toumatou), bracken	Major
Forest	Southern beech on river flats and along margins of the IBA.	Minor

Black-billed Gulls, and to a lesser extent Black-fronted Terns will feed on adjacent pasture, especially during or immediately following ploughing. They will also hawk flying insects over pasture, riverbeds and even forested areas.



Threats to the site (pressure) - IUCN Classifications

Threat Level	Notes	Timing	Scope	Severity
Invasive & other prob- lematic species, genes & diseases	Invasive alien species - mustelids, cats, rats, hedgehogs	Happening now	Some of area/ population (10-49%)	Moderate to rapid deterioration
Human intrusions and disturbance	Recreational activities by river users - includes dogs.	Happening now	Some of area/ population (10-49%)	Slow but significant deterioration
Natural system modifications	Water management/use - abstraction of ground water (agricultural use)	Happening now	Some of area/ population (10-49%)	Slow but significant deterioration
Natural system modifications	Weed encroachment on nesting sites (rivers)	Happening now	Some of area/ population (10-49%)	Slow but significant deterioration
Climate change and severe weather	Floods	Likely in short term (within 4 years)	Some of area/ population (10-49%)	Moderate to rapid deterioration

#### **References:**

McClellan, R. 2008. The Ecology and Management of Southland's Black-billed Gulls. Thesis for the degree of Doctor of Philosophy, University of Otago, Dunedin. 253 pp.

O'Donnell, C.F.J., Hoare, J.M. 2011. Meta-analysis of status and trends in breeding populations of black-fronted terns (Chlidonias albostriatus) 1962-2008. New Zealand Journal of Ecology 35: 32-43

Robertson, C.J.R., Hyvönen, P., Fraser, M.J., Pickard, C.R. 2007. Atlas of bird distribution in New Zealand 1999-2004. Ornithological Society of New Zealand, Inc. Wellington.



